QE
756
J29
N53

日本古生物学會報告・紀事

Transactions and Proceedings of the Palaeontological Society of Japan

New Series

No. 81～No. 88

1971–1972

日本古生物学会

Palaeontological Society of Japan

The heading in Japanese commemorates the handwriting of Prof. Matajiro YOKOYAMA, father of Japanese palaeontology, who was a professor of stratigraphy and palaeontology at the Geological Institute, Imperial University of Tokyo.

Fossil on the cover is left lower M2 of *Palaeoloxodon naumanni* (MAKIYAMA, 1924) from the uppermost part of the Tokyo formation (Upper Pleistocene) at Ikebukuro, Tokyo.

CONTENTS

Number 81 (Published April 20, 1971)

Transactions

Article		Page	Plate
574.	KATO, Makoto: J. FLEMING's species of British Lower Carboniferous corals	1–10	1
575.	TAKAHASHI, Kiyoshi: Microfossils from the Pleistocene sediments of the Ariake Sea area, west Kyushu	11–26	2–5
576.	NODA, Hiroshi: New anadarid and associated molluscan fauna from the Haneji Formation, Okinawa-jima, Ryukyu Islands	27–51	6–7
577.	HATAI, Kotora and KOTAKA, Tamio: Some coprolites from Wakayama Prefecture	52–58	
	Proceedings	59–60	

Number 82 (Published June 30, 1971)

Transactions

578.	HU, Chung-Hung and TAN, Li-Lin: Ontogenies of two Upper Cambrian trilobites from northern Black Hills, South Dakota	61–72	8–9
579.	HAYAMI, Tomoko: Some Neogene Cheilostomata (Bryozoa) from Okinawa-jima	73–92	10–12
580.	MURATA, Masafumi: Fusulinid biostratigraphy and molluscan fauna from the uppermost part of the Sakamotozawa formation, and the pre-Kanokura unconformity, in the southern part of the Kitakami massif, Northeast Japan	93–116	13–14
581.	BURCKLE, Lloyd H.: Correlation of Late Cenozoic marine sections in Japan and the Equatorial Pacific	117–128	

Number 83 (Published September 20, 1971)

Transactions

582.	KURIHARA, Kenji: Foraminifera from the Hayama Group, Miura Peninsula	131–142	15

583. NISHIDA, Shiro: Nannofossils from Japan IV. Calcareous nannoplankton fossils from the Tônohama Group, Shikoku, Southwest Japan 143-161 16-18
584. HATAI, Kotora and NODA, Hiroshi: Peculiar markings on a sandstone layer of the Hagino Formation, Nagano Prefecture.....162-165
585. MASUDA, Kôichirô: On some *Patinopecten* from North America 166-178 19-21

Number 84 (Published December 30, 1971)

Transactions

586. MATSUMARU, Kuniteru: The genera *Nephrolepidina* and *Eulepidina* from New Zealand 179-189 22-23
587. KIMURA, Tatsuaki and SEKIDO, Shinji: The discovery of the cycad-like leaflets with toothed margin from the Lower Cretaceous Itoshiro Sub-group, the Tetori Group, Central Honshu, Japan....190-195 24
588. HAYAMI, Tomoko: On some Bryozoa from near Namioka-cho, Minami-Tsugaru-gun, Aomori Prefecture, Japan 196-204
589. MASUDA, Kôichirô: *Amussiopecten* from North America and northern South America.....205-224 25-26
590. TASHIRO, Masayuki: Upper Cretaceous glycrmerids in Japan ..225-242 27-28
Proceedings 243-244

Number 85 (Published April 20, 1972)

Transactions

591. HU, Chung-Hung: Ontogeny of three *Cedaria* zone trilobites from Upper Cambrian, Montana 245-259 29-30
592. YAMAGIWA, Nobuo and SAKA, Yukiyasu: On the *Lepidolina* zone discovered from the Shima Peninsula, Southwest Japan 260-274 31-32
593. SAKAGAMI, Sumio: The Triassic Bryozoa from Kusaka, Sakawa basin, Shikoku, Japan 275-279 33
594. SATO, Tadashi: Some Bajocian ammonites from Kitakami Northeast Japan 280-292 34
Proceedings 293

Number 86 (Published June 30, 1972)

Transactions

595. FUJI, Norio: Fossil spores and pollen grains from the Neogene deposits in Noto Peninsula, Central Japan-IV. A palynological study of the Late Miocene Tsukada Member 295-318 35-38
596. HATAI, Kotora and NODA, Hiroshi: A Problematica from the Mizuho-To of Niigata Prefecture.....319-324 39
597. TASHIRO, Masayuki: On the surface ornamentation of the pennatae trigoniids, and on three new species of the trigoniids from the Himenoura Group, Kyushu, Japan 325-339 40-41
598. KOIZUMI, Itaru: Marine diatom flora of the Pliocene Tatsunokuchi-Formation in Fukushima Prefecture 340-359 41-43
599. KIMURA, Tatsuaki and SEKIDO, Shinji: *Ctenis* species from the Itoshiro Sub-group (Lower Cretaceous), the Tetori Group, Central Honshu, Japan 360-368 44-45
600. CHOI, Dong Ryong: *Colania douvillei* (OZAWA), a fusulinid Foraminifera, from the Northern Kitakami Mountains, NE Japan....369-374 46

Number 87 (Published September 30, 1972)

Transactions

601. MATSUMOTO, Tatsuro, MURAMOTO, Tatsuo and INOMA, Akitoshi: Two small desmoceratid ammonites from Hokkaido (Studies of the Cretaceous ammonites from Hokkaido and Saghalien-XXIV)..377-394 47
602. MASUDA, Koichiro: *Swiftopecten* of the Northern Pacific 395-408 48-49
603. HATAI, Kotora, NODA, Hiroshi and OGASAHARA, Kenshiro: Scaphopoda-like fossils from the Udo formation (Miocene) of Miyazaki Prefecture, Japan 409-413
604. OKIMURA, Yuji: Permo-Carboniferous endothyraceans from Japan, Part 1. Biseriamminidae 414-428 50-51
605. TAKAYAMA, Toshiaki: A note on the distribution of *Braarudosphaera bigelowi* (GRAN & BRAARUD) DEFLANDRE in the bottom sediments of Sendai Bay, Japan 429-435
606. SADA, Kimiyoshi: Fusulinids of the *Profusulinella* zone of the Taishaku limestone. (Studies of the stratigraphy and microfossil faunas of the Carboniferous and Permian Taishaku Limestone in

West Japan, No. 2) 436-445 52-53
Proceedings 446

Number 88 (Published December 20, 1972)

Transactions

607. ISHIBASHI, Takeshi: Upper Triassic cephalopods from the Tanoura district, Kumamoto Prefecture, Japan 447-457 54
608. HATAI, Kotora and NODA, Hiroshi: A fossil worm trail from the Upper Cretaceous Nakaminato formation, Ibaraki Prefecture 458-461 55
609. NODA, Masayuki: Ammonites from the Mesozoic Yamabu formation, Kyushu 461-471 56
610. NODA, Hiroshi: Some fossil Pteropoda from Miyazaki and Okinawa Prefectures, Southwest Japan 472-484 57
Proceedings 485
Systematic Index 486-488
Index of Families, Genera and Species .. 489-492